

# Assimilating Satellite-derived Soil Moisture and Ingesting Real-time Vegetation into WRF-Hydro using the NASA LIS

**Nicholas Elmer**<sup>1,3</sup>, Clay Blankenship<sup>2,5</sup>, Jonathan Case<sup>3,5</sup>,  
Bradley Zavodsky<sup>4,5</sup>, and Andrew Molthan<sup>4,5</sup>

<sup>1</sup>Department of Atmospheric Science, University of Alabama in Huntsville, Huntsville, Ala.

<sup>2</sup>Universities Space Research Association (USRA) Space Technology, Institute, Huntsville, Ala.

<sup>3</sup>ENSCO, Inc., Huntsville, Ala.

<sup>4</sup>Earth Science Office, NASA MSFC, Huntsville, Ala.

<sup>5</sup>NASA Short-term Prediction Research and Transition (SPoRT) Center, Huntsville, Ala.

Collaborators:

Joseph Santanello (NASA GSFC)

David Gochis (NCAR)




**January 24, 2017**

**31<sup>st</sup> Conference on Hydrology**

**97<sup>th</sup> AMS Annual Meeting, Seattle, Wash.**

# Operational SPoRT LIS




- CONUS, 3-km resolution
- NASA LIS (Kumar et al. 2006; Peters-Lidard et al. 2007) used to perform long-term integration of Noah Land Surface Model (LSM) updated in real-time
- Assimilation of soil moisture gives even more accurate LSM soil moisture fields
  - Currently undergoing operational assessment
- Output used for situational awareness and local modeling by forecasters at select NWS offices and international forecasting agencies



# Application: Areal Flood Potential

*March*


Moderate antecedent soil moisture



Moderate-heavy precipitation




Moderate river flooding and numerous flooding reports





*September*


Low antecedent soil moisture



Heavy precipitation




Isolated minor flooding




- Contrasting antecedent soil moisture likely played a strong role in the different outcomes
- Local, subjective analysis of several events suggests typical moderate-heavy synoptic rainfall events over deep-layer relative soil moisture values exceeding 55-60% will lead to more substantial moderate or heavier flooding events




# Assimilation of Soil Moisture Data





*Model soil moisture concentration forced only by precipitation and misses magnitude of irrigation-saturated MS Valley*



*SMOS observes irrigated fields*




*Blended analysis of model and observations better represent irrigated area and should result in improved weather and hydrologic modeling*




# SMOS DA Validation

0–10 cm Change in Correlation



Degraded w/ DA

Root Zone Change in Correlation



Improved w/DA


Variable	0-10 cm Soil Moisture				
# Stations	194				
Experiment	OPL	NOBC	BC1	BCS	BCV
Bias	<b>-0.000</b> $\pm$ 0.011	-0.026 $\pm$ 0.011	-0.023 $\pm$ 0.011	-0.005 $\pm$ 0.011	-0.025 $\pm$ 0.011
RMSE	<b>0.082</b> $\pm$ 0.005	0.087 $\pm$ 0.006	0.086 $\pm$ 0.005	<b>0.082</b> $\pm$ 0.005	0.087 $\pm$ 0.006
Unbiased RMSE	0.046 $\pm$ 0.003	<b>0.043</b> $\pm$ 0.002	<b>0.043</b> $\pm$ 0.002	0.044 $\pm$ 0.003	<b>0.043</b> $\pm$ 0.002
Correlation	0.451 $\pm$ 0.023	<b>0.573</b> $\pm$ 0.027	0.569 $\pm$ 0.026	0.539 $\pm$ 0.025	0.561 $\pm$ 0.026

Assimilation of SMOS using soil classification bias correction results in best overall configuration for bias, RMSE, and  $r^2$



# WRF-Hydro System

- Weather Research and Forecasting model hydrological extension package (WRF-Hydro; Gochis et al. 2013)
- Extensible, high-resolution hydrologic routing and streamflow modeling framework
- Contains column land surface, terrain routing, and channel routing modules
- National Water Model (NWM; Office of Water Prediction 2017) runs instantiation of WRF-Hydro operationally



(NCAR 2017)

# Coupling LIS and WRF-Hydro

- Collaborative project between NASA GSFC and NCAR (Santanello et al. 2015)
  - Funded by NASA's Modeling, Analysis, and Prediction (MAP) program
  - Couple LIS and WRF-Hydro in the Earth Science Modeling Framework (ESMF), which will enable operational linking of these two systems
- Plan to leverage this project to assimilate/integrate NASA mission datasets in WRF-Hydro using the LIS Ensemble Kalman Filter (EnKF)




# Assimilation of SMAP/SMOS into WRF-Hydro

- Soil moisture (**SMAP/SMOS**)
- Snow cover (MODIS/VIIRS) and snow water equivalent (AMSR2)
- Total terrestrial/ground water (GRACE/GRACE-Follow On)
- Other future NASA missions (ICESat-2, Landsat-9)

- Inundation and Streamflow (NISAR, SWOT)


## NASA LIS

Column Land Surface Models:




Output Variables:  
Evapotranspiration  
Soil moisture/Soil Ice  
Snowpack/snowmelt  
Runoff  
Radiation Exchange  
Energy Fluxes  
Plant Water Stress

## Channel & Reservoir Routing Models: Hydrologic and Hydraulic



Output Variables:  
Streamflow  
River Stage  
Flow Velocity  
Reservoir Storage  
& Discharge

2-way coupling



Terrain Routing Models:  
Overland, subsurface flow

Output Variables:  
Stream Inflow, Surface Water Depth, Groundwater Depth, Soil Moisture

1-way coupling or  
2-way coupling




WRF-Hydro  
System



# Evaluating LIS fields in WRF-Hydro

## December 2015 Alabama Flood


2015-12-25 1200 UTC




- Noah-MP initialized with SPORT-LIS soil moisture, soil temperature, surface skin temperature, and vegetation fraction
- Multi-Radar Multi-Sensor (MRMS) 1-hr gauge corrected accumulated precipitation (background field;  $mm\ hr^{-1}$ )
- “Cold start” of hydrological model (i.e., streambeds initially dry)

# Impacts of Real-time Vegetation

Monthly Mean LAI - December




Real-time LAI (VIIRS/LIS) - 20151223



- Ongoing project to quantify impact of VIIRS real-time vegetation on simulated soil moisture and streamflow
- Larger deviations from climatology also likely in early spring and late fall

# Impacts of Real-time Vegetation

- 20-29 December 2015 modeled streamflow
- climatological GVF (dashed black line) and real-time VIIRS GVF (solid green/red line)
- Absolute difference (gray solid line)
- Replacing climatological GVF with real-time GVF results in approximately 1% change in streamflow (i.e., minimal impact for high-flow event)
- Greater differences expected for low-flow events



# LIS as Assimilation Framework for NWM

- Currently, the NWM does not have a system for assimilating land surface satellite observations
- The LIS system is a strong candidate given both the long history of the LIS and its linkage through the ESMF
- LIS EnKF enables assimilation of satellite-based observations (e.g., SMAP, SMOS, MODIS, VIIRS, AMSR-2, ICESat-2, etc.)
- SPoRT-LIS is being upgraded to include a Noah-MP LSM run using NWM configurations to demonstrate impacts of NASA datasets in NWM





# Summary and Future Work

- SPoRT is assimilating/ingesting satellite soil moisture and vegetation measurements into the operational SPoRT-LIS
- NASA LIS being coupled to WRF-Hydro by GSFC and NCAR
- In collaboration with the National Water Center (NWC), SPoRT is developing an offline, experimental version of the NWM to evaluate the impact of current and future NASA mission datasets (e.g., SMAP soil moisture, VIIRS real-time vegetation, SWOT surface water elevations)



Nicholas Elmer  
[nicholas.j.elmer@nasa.gov](mailto:nicholas.j.elmer@nasa.gov)

## NASA SPoRT

Webpage: <http://weather.msfc.nasa.gov/sport/>

Blog: <https://nasasport.wordpress.com/>

Facebook: NASA SPoRT Center

Twitter: @NASA\_SPoRT

### References

- Blankenship, C. B., J. L. Case, B. T. Zavodsky, and W. L. Crosson, 2016: Assimilation of SMOS Retrievals in the Land Information System. *IEEE Trans. Geo. Rem. Sens.*, **54**(11), 6320-6332, doi:10.1109/TGRS.2016.2579604.
- Elmer, N., B. Zavodsky, A. Molthan, J. Case, and C. Blankenship, 2016: Impacts of real-time satellite-derived vegetation on WRF-Hydro simulated streamflow. Abstract H51H-1618 presented at 2016 Fall Meeting, AGU, San Francisco, Calif., 12-16 Dec.
- Gochis, D., W. Yu, and D. Yates, 2013: The NCAR WRF-Hydro technical description and user's guide: version 1.0, 120 pp., [http://www.ral.ucar.edu/projects/wrf\\_hydro/images/WRF\\_Hydro\\_Technical\\_Description\\_and%20User\\_Guide\\_v1.0.pdf](http://www.ral.ucar.edu/projects/wrf_hydro/images/WRF_Hydro_Technical_Description_and%20User_Guide_v1.0.pdf).
- Kumar, S. V., et al., 2006: Land Information System - an interoperable framework for high resolution land surface modeling. *Environmental Modeling & Software*, **21**(10), 1402-1415, doi:10.1016/j.envsoft.2005.07.004.
- NCAR, 2017: WRF-Hydro Modeling System. Research Applications Library, [https://www.ral.ucar.edu/sites/default/files/public/projects/wrf\\_hydro/](https://www.ral.ucar.edu/sites/default/files/public/projects/wrf_hydro/).
- Office of Water Prediction, cited 2017: The National Water Model. Office of Water Prediction, <http://water.noaa.gov/about/nwm>.
- Peters-Lidard, C. D., et al., 2007: High-performance Earth system modeling with NASA/GSFC's Land Information System. *Innovations Syst. Softw. Eng.*, **3**(3), 157-165, doi:10.1007/s11334-007-0028-x.
- Santanello, Jr., J. A., S. V. Kumar, D. Gochis, C. DeLuca, C. D. Peters-Lidard, 2015: An integrated, observation-driven hydrological modeling system using LIS and WRF-Hydro enabled by ESMF. 2015 NASA Research Opportunities in Space and Earth Science Funded Proposal (NNH15ZDA001N-MAP), 21 pp.

